

Quantitative Determination of a Microscopic Mechanism of High T_c Superconductivity

Eugene Demler and Shou-Cheng Zhang

Department of Physics, Stanford University Stanford, CA 94305

(February 1, 2008)

A major challenge in theoretical attempts to understand the microscopic mechanism of high T_c superconductivity is to quantitatively account for the superconducting condensation energy [1–6]. Microscopic model commonly used to describe the superconducting copper-oxides, the t - J model [7], is believed to capture the essential physics of these materials: the interplay of electrons kinetic energy and their antiferromagnetic exchange interaction. Within the t - J model, the condensation energy can be related to the change in the dynamical spin structure factor between the superconducting and the normal states [8]. By analyzing previous experimental data, we show that the change associated with the resonant neutron scattering peak [9–11] found in $YBa_2Cu_3O_7$ can quantitatively account for the condensation energy. We argue that this analysis suggests a microscopic mechanism for high T_c superconductivity, where antiferromagnetic exchange energy is saved in the superconducting state through the coupling to a particle-particle resonance [12–16].

One reason for the absence of a generally accepted microscopic explanation of high T_c superconductivity (HTSC) is the lack of a quantitatively precise definition of a “mechanism”. In the traditional BCS [1] superconductors, the phonon mechanism is quantitatively verified by the isotope substitution and single particle tunneling experiments. It is a widely held view that the HTSC involve strong electronic interaction, while phonon mediated interaction is not the major driving force for superconductivity. To date, many different “mechanisms” have been proposed, but very few quantitatively falsifiable predictions have been checked against experimental data.

A system undergoes a transition from the normal to the superconducting state because it can lower the total free energy. The condensation energy E_C , defined as the energy difference between the normal state, extrapolated to zero temperature, (E_N) and the superconducting state (E_S), can be directly measured through the thermodynamical critical field H_c by the following relation [1]:

$$E_C = E_N - E_S = \frac{H_c^2}{8\pi} V_0 \quad (1)$$

Here we choose $V_0 = a \times b \times c$ to be the volume of a unit cell (in the high T_c cuprates $a = b$ with reasonably good accuracy) and define the energies inside one unit cell. Alternatively, the condensation energy can also be found by integrating the difference in specific heats in the (extrapolated) normal and superconducting states from $T = 0$ to the superconducting transition temperature T_c [2,3].

On the other hand, the condensation energy can be computed in principle from a microscopic Hamiltonian. Ideally, one would like to start from the full Hamiltonian involving electronic kinetic energy, the Coulomb interaction energy and the electron lattice coupling energy. In fact Chester [4] studied condensation energy of such a Hamiltonian and showed how various contributions to the condensation energy could in principle be determined experimentally. Among many theories of HTSC, only the interlayer tunneling mechanism [5] made a quantitatively falsifiable prediction about the condensation energy. More recently, Leggett showed that the change in the interlayer Coulomb energy can be directly related to the frequency and momentum integral of the dynamic charge susceptibility, a quantity which can be measured experimentally [6]. He argued that a microscopic theory of HTSC should quantitatively predict the frequency and the momentum range at which the change of the dynamical susceptibility occurs, and quantitatively account for the condensation energy.

However, the energy scale involved in HTSC is much smaller than the Coulomb interaction. In order to understand the mechanism for HTSC, it is useful to start with an effective Hamiltonian which correctly describes the basic physics below the Coulomb energy scale. Recently, Scalapino and White [8] initiated the investigation of condensation energy in the

t - J model, which is defined by

$$\mathcal{H} = -t \sum_{\langle ij \rangle} c_{i\sigma}^\dagger c_{j\sigma} + J \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j \quad (2)$$

Here $c_{i\sigma}^\dagger$ is electron creation operator on site i with spin σ , $\mathbf{S}_i$ is the electron spin operator, $\langle ij \rangle$ denotes a pair of near neighbor sites and the Hamiltonian acts on the space of no doubly occupied sites. Equation (2) describes the Hamiltonian of one layer only. In a bi-layer system like $YBa_2Cu_3O_{6+x}$, there are two layers per unit cell, with additional antiferromagnetic (AF) coupling $J_\perp$ between the layers. If a system described by this Hamiltonian undergoes a superconducting transition away from half-filling, the kinetic energy E_t is expected to increase slightly, but the decrease in the exchange energy E_J may be significantly larger, so that a superconducting ground state is realized.

In a tight binding model, the mean kinetic energy E_t can be expressed as a frequency integral of the optical conductivity $\sigma(\omega)$ [17–19]. Then the contribution of E_t to the condensation energy can be measured by comparing $\sigma(\omega)$ in the normal and the superconducting states. In the low frequency regime, the Drude peak in the normal state collapses into a δ function peak in the superconducting state, usually with conserved weight [20]. Therefore, the change in the kinetic energy has to be determined from $\sigma(\omega)$ in the higher frequency range.

Scalapino and White [8] made a insightful observation that the change in the J term in equation (2) can also be directly expressed as a frequency and momentum integral of the dynamic spin structure factor $S(q, \omega)$. For bi-layer materials such as $YBa_2Cu_3O_{6+x}$ it is convenient to separate the even and odd parts of $S(q, \omega)$ with respect to bi-layer interchange within a unit cell

$$S(q, \omega) = S^{even}(q, \omega) \cos^2\left(\frac{q_z d_{Cu}}{2}\right) + S^{odd}(q, \omega) \sin^2\left(\frac{q_z d_{Cu}}{2}\right) \quad (3)$$

Here d_{Cu} is the distance between nearest-neighbor Cu - O planes, so S^{even} and S^{odd} describe the in-phase and out-of-phase spin fluctuations in the two planes constituting one bi-layer. Then $\Delta E_J = E_J^N - E_J^S$ may be obtained as (per unit cell)

$$\Delta E_J = \frac{3}{2} \left(\frac{a}{2\pi} \right)^2 \int d^2 q \int \frac{d(\hbar\omega)}{\pi} \{ [S_N^{even}(q, \omega) - S_S^{even}(q, \omega)] \left[\frac{1}{2} J(\cos(q_x a) + \cos(q_y a)) + \frac{1}{4} J_\perp \right] + [S_N^{odd}(q, \omega) - S_S^{odd}(q, \omega)] \left[\frac{1}{2} J(\cos(q_x a) + \cos(q_y a)) - \frac{1}{4} J_\perp \right] \} \quad (4)$$

From [21,22] it is known that $J_\perp$ is much smaller than J , so the $J_\perp$ term in equation (4) may be safely neglected. Scalapino and White pointed out [8] that since $S(q, \omega)$ can be measured directly in neutron scattering experiments, ΔE_J can therefore in principle be measured. If ΔE_J turns out to be much smaller than the experimentally measured condensation energy E_c , this gives a convincing way to rule out any mechanism of high T_c superconductivity based on the $t - J$ model. On the other hand, if ΔE_J turns out to be greater than E_c , one would have identified a important driving force for superconductivity, and can quantitatively test any theoretical mechanism of HTSC based on the $t - J$ model, which should predict where in frequency and momentum space the change of $S(q, \omega)$ occurs.

However, one should apply this line of reasoning with extreme care. The quantity $S_N(q, \omega)$ in equation (4) is not the normal state spin structure factor above T_c , but rather an extrapolated normal state quantity at $T = 0$. Experimentally, one has to carefully identify features in $S(q, \omega)$ which changes abruptly at T_c . Theoretically, one has to identify contributions to $S(q, \omega)$ which are absent in the normal state and only present in the superconducting state, so that the difference in equation (4) can be rigorously defined. Recently, a resonant neutron peak has been discovered in the family of $YBa_2Cu_3O_{6+x}$ superconductors [9–11]. In the optimally doped $YBa_2Cu_3O_7$ superconductor with $T_c = 91K$, the resonance peak occurs in the spin scattering channel at energy $\omega_0 = 41 \text{ meV}$. It is centered around momentum $Q = (\frac{\pi}{a}, \frac{\pi}{a})$ and it occurs in the odd channel with respect to bi-layer interchange. Perhaps the most remarkable feature of the resonance peak is its onset at the superconducting transition temperature T_c . Above T_c , there is no experimentally identifiable spectral weight in this frequency and momentum range. Therefore, this experiment is ideally suited for analyzing its contribution to the condensation energy.

We constructed a theory [12] of the neutron resonance peak by first identifying a resonance in the spin-triplet particle-particle (p-p) channel defined by the following operator

$$\pi^\dagger = \sum_k (\cos k_x - \cos k_y) c_{k+Q\uparrow}^\dagger c_{-k\uparrow}^\dagger \quad (5)$$

In a series of exact diagonalization studies on the Hubbard and t - J models, Meixner, Hanke and the two of us [13] and Eder, Hanke and one of us (SCZ) [14] demonstrated the existence of such a triplet p-p resonance for all doping range. A triplet p-p resonance can not be detected in the neutron scattering experiments on the normal state. However, below T_c , the d-wave superconducting order parameter mixes a p-p resonance into the dynamical spin structure factor, leading to a sharply defined collective mode in the triplet particle-hole (p-h) channel (the so-called π -resonance). The contribution from the p-p resonance to the p-h channel below T_c is given [12,15] by

$$\int d(\hbar\omega) S(q, \omega) = \frac{2|\langle\Delta_d\rangle|^2}{1-n} \quad (6)$$

where $\langle\Delta_d\rangle$ is the dimensionless d -wave superconducting order parameter (not the energy gap) and n is the density of electrons. This simple estimate of the spectral weight of the π resonance at $q = Q$ agrees reasonably well with the absolute intensity measured by Fong *et al.* [11]. On the other hand, the width of the π resonance in momentum space can not be calculated reliably. Within the $SO(5)$ theory [16], the π resonance has been interpreted as a pseudo-Goldstone boson of the spontaneous $SO(5)$ symmetry breaking in the superconducting state. Its existence is therefore directly correlated with the existence of the superconducting long range order.

Since the coupling to a p-p resonance is only possible in the superconducting state, it indeed contributes to the difference in $S(q, \omega)$ as required by equation (4). In this work we wish to point out that the logic leading to our explanation of the π resonance below T_c can be reversed to provide a microscopic mechanism of HTSC. On one hand, the existence of the π resonance is a unique property of the superconducting state, on the other hand, its existence around Q lowers E_J , as we see from equation (4). Therefore, the system undergoes a superconducting transition, so that a π resonance mode can emerge, which in turn lowers E_J . In other words, no matter how antiferromagnetically correlated the normal state is, a superconducting state can always further lower the exchange energy by coupling a p-p

resonance into the dynamic spin structure factor $S(q, \omega)$. Our mechanism for HTSC makes a quantitatively precise prediction which can be checked experimentally. Once we assume that the HTSC is dominantly driven by the emergence of a new collective mode, the π resonance, this mechanism predicts that the condensation energy is determined by the zero temperature spectral weight of the π resonance, and the energy saving occurs precisely in the frequency and momentum range of the π resonance.

Since our mechanism predicts an additional spectral weight around $q = Q$ in the superconducting state, a careful reader may wonder what happens to the spectral sum rule, which states that $S(q, \omega)$ integrated over both q and ω (without the $f(q) = \cos(q_x a) + \cos(q_y b)$ factor in equation (4)) should be a constant, independent of the nature of the ground state. Of course our additional spectral weight should be compensated by a depletion of spectral weights from other parts in the momentum space. However, since $f(q)$ has a absolute minimum at $q = Q$, their contributions to (4) can not completely cancel the π resonance contribution. In fact, depletion of the spectral weight near $q = 0$ would lead to additional enhancement of the condensation energy.

Since both the condensation energy and the spectral weight of the neutron resonance peak have been measured experimentally, it is straightforward to check this prediction. Within Landau-Ginzburg theory, $H_c = \frac{\Phi_0}{2\sqrt{2}\pi\xi\lambda}$, where $\Phi_0 = \frac{hc}{2e}$ is the superconducting flux quantum, ξ is the coherence length and λ is the London penetration depth. In $YBa_2Cu_3O_7$, these two length scales are determined to be in the range of $\xi = 12 - 20 \text{ \AA}$ and $\lambda = 1300 - 1500 \text{ \AA}$. Using $a = b = 3.85 \text{ \AA}$ and $c = 11.63 \text{ \AA}$ this gives a condensation energy of $E_C = 3.3 - 12 \text{ K}$ per unit cell. On the other hand, E_C can also be measured directly in specific heat experiments. Loram et al. reported (see page 251 of [2] and page 247 of [3]) a value of $E_C = 6 \text{ K}$ per unit cell for $YBa_2Cu_3O_7$. Let us now turn to the experimental measurement of ΔE_J given by equation (4). In practice, it is very hard to measure the change of $S(q, \omega)$ throughout the frequency and momentum range. However, since our mechanism predicts that the major part of the condensation energy is saved in the range of $\omega_0 = 41 \text{ meV}$ and a 2D momentum

transfer $Q = (\frac{\pi}{a}, \frac{\pi}{a})$, we can analyze the neutron scattering data by assuming that the contributions of the other parts do not change significantly at T_c . Indeed, for frequencies above the resonance the neutron scattering intensity does not change as the system goes from normal to superconducting. For frequencies below and in the resonance range the normal state intensity $S_N(q, \omega)$ is below experimental sensitivity limit in both even and odd channels. Because of the sum rule discussed above we may not assume that it vanishes identically, so we make an assumption that in this frequency range the spectral weight $S_N(q, \omega)$ is spread uniformly in momentum space and therefore does not contribute to (4). Below T_c , a neutron resonance peak emerges in the odd channel, and $S_S^{odd}(q, \omega)$ has been measured in absolute units in the range of the resonance. The dimensionless quantity $\int d(\hbar\omega) S_S^{odd}(Q, \omega)$ is measured to be 0.52 at $T = 10\text{ K}$ by Fong *et al.* [11] (Theoretical estimate based on references [12,15] gives 0.32 for this value). The resonance has a Lorentzian profile centered at $q = Q$ with a width $\kappa_{2D} = 0.23\text{\AA}^{-1}$ [11], so the 2D momentum integral can be easily estimated.

$$\Delta E_J = \frac{3}{2} \pi \left(\frac{a}{2\pi} \kappa_{2D} \right)^2 \times \frac{1}{2} \times \frac{0.52}{\pi} \times 2 \times J = 0.016J \quad (7)$$

Taking $J = 100\text{ meV}$ we obtain $\Delta E_J = 18\text{ K}$. The actual condensation energy should be smaller than ΔE_J since the kinetic energy is expected to increase below the superconducting state. Taking this into account, we see that the spectral weight of the π resonance peak can quantitatively account for the condensation energy measured from H_c and specific heat experiments, in reasonable agreement with the prediction based on our mechanism.

It is worthwhile to compare our mechanism for HTSC with the spin fluctuation pairing mechanism [23–25]. All these mechanisms are based on the AF exchange interaction interaction E_J . Our analysis of the neutron data confirms that a large part of the condensation energy indeed arises from the AF exchange interaction [8]. However, our mechanism differs from spin fluctuation pairing mechanism, both conceptually and quantitatively. The idea of spin fluctuation pairing is directly borrowed from the phonon mediated pairing in the traditional BCS superconductors. It requires sizable AF spin fluctuations in the normal state, in

order to pair the fermi liquid like quasiparticles. In contrast, in our mechanism, the AF spin fluctuation does not play an important role in driving superconductivity, and is neglected in the normal state of optimally doped $YBa_2Cu_3O_7$ superconductor. In the superconducting state, the π resonance can be viewed as a kind of spin fluctuation in a particular frequency and momentum range. From the point of view of the $SO(5)$ theory [16], the superconducting state is obtained from the AF state by a rotation, it is therefore natural to expect it to have more AF correlation than the normal state. Therefore, the crucial conceptual difference between these two mechanisms lies in the fact that normal state AF spin fluctuations are not required in our mechanism. This conceptual difference has a direct quantitative consequence. From equation (4) we see that the presence of spin fluctuation spectral weight in the normal state contributes negatively to the condensation energy. It is not clear why the spin fluctuation model would predict a change in the frequency integrated weight of $S(q, \omega)$ near $q = Q$. In fact, most theories of the neutron resonance peak based on the spin fluctuation models [26,27] assume a preexisting overdamped spin fluctuation mode in the normal state and only predict a reduction of damping below T_c , with essentially conserved weight.

Our mechanism provides a natural explanation to the doping dependence of the condensation energy. We attribute the condensation energy to the difference in exchange energies in superconducting and normal states. More underdoped materials have considerably more AF correlations in the normal state, as known from the neutron scattering [28]. Although in the superconducting state the weight of the resonance is enhanced [29], the difference between the normal and the superconducting exchange energies becomes smaller. This is consistent with the results of Loram *et al.* who find a decrease in the condensation energy with decreased doping in $Y_{0.8}Ca_{0.2}Ba_2Cu_3O_{6+x}$ [30]. Until this point we have been discussing the relation between the condensation energy and the resonant peaks in neutron scattering at zero temperature. It would be interesting to see whether this idea works at finite temperature as well. Without going into any details we would like to point at a striking similarity between the temperature dependencies of H_c [2] and the resonance intensity [29,31] for the underdoped cuprates. Not only both quantities scale similarly below T_c , but

they both have tails extending above T_c , presumably arising from significant pairing fluctuations in the pseudogap regime of the cuprates. We postpone the quantitative analysis of these phenomena to future investigations.

We would like to thank P. Dai, H. Fong, S. Heyden, B. Keimer, H. Mook, and D. Scalapino for useful discussions.

-
- [1] Shrieffer, J.R. Theory of Superconductivity, *Addison-Wesley, MA*, (1964).
 - [2] Loram, J. *et al.* The electronic specific heat of $YBa_2(Cu_{1-x}Zn_x)_3O_7$ from 1.6 to 300K. *Physica*, **C 171** 243-256 (1990).
 - [3] Loram, J. *et al.* Electronic specific heat of $YBa_2Cu_3O_{6+x}$ from 1.8 to 300K. *Journal of Superconductivity*, **7** 243-249 (1994).
 - [4] Chester, G.V. Difference between normal and superconducting states of a metal. *Phys. Rev.*, **103** 1693-1699 (1956).
 - [5] Chakravarty, S. *et al.* Interlayer tunneling and gap anisotropy in high temperature superconductors. *Science*, **261** 337-340 (1993).
 - [6] Leggett, A. Where is the energy saved in cuprate superconductivity? *preprint*.
 - [7] Zhang, F.C. and Rice, T.M., Effective Hamiltonian for the superconducting copper-oxides. *Phys. Rev. B*, **37**, 3759 (1988)
 - [8] Scalapino, D.J. and White, S., The superconducting condensation energy and an antiferromagnetic exchange based pairing mechanism. *preprint*, cond-mat/9805075.
 - [9] Mook, H. *et al.* Polarized neutron determination of the magnetic excitation in $YBa_2Cu_3O_7$. *Phys. Rev. Lett.*, **70** 3490-3493 (1993).
 - [10] Fong, H.F. *et al.* Phonon and magnetic neutron scattering at 41 meV in $YBa_2Cu_3O_7$. *Phys.*

- Rev. Lett.*, **75** 316-319 (1995).
- [11] Fong, H.F. *et al.* Polarized and unpolarized neutron scattering study of the dynamic spin susceptibility in $YBa_2Cu_3O_7$. *Phys. Rev. B*, **54** 6708-6720 (1996).
 - [12] Demler, E. and Zhang, S.C. Theory of resonant neutron scattering of high T_c superconductors. *Phys. Rev. Lett.* **76** 4126-4129 (1995).
 - [13] Meixner, S. *et al.* Finite size studies on the $SO(5)$ symmetry of Hubbard model. *Phys. Rev. Lett.*, **79** 4902-4905 (1997).
 - [14] Eder, R., Hanke, W., and Zhang, S.C. Numerical evidence for $SO(5)$ symmetry and superspin multiplets in the two dimensional $t - J$ model. *Phys. Rev. B.*, **57** 13781-13789 (1998).
 - [15] Demler, E., Kohno, H. and Zhang, S.C. π excitation of the $t - J$ model. *preprint*, cond-mat/9710139.
 - [16] Zhang, S.C. A unified theory based on $SO(5)$ symmetry of superconductivity and antiferromagnetism. *Science*, **275** 1089-1096 (1997).
 - [17] Shastry, S. and Sutherland, B. Twisted boundary conditions and effective mass in Heisenberg-Ising and Hubbard rings. *Phys.Rev.Lett.*, **65** 243-246 (1990).
 - [18] Scalapino, D., White, S. and Zhang, S.C. Superfluid density and the Drude weight of the Hubbard model. *Phys.Rev.Lett.*, **68** 2830-2833 (1992).
 - [19] Chakravarty, S. Do electrons change their c-axis kinetic energy upon entering the superconducting state? *preprint*, cond-mat/9801025.
 - [20] Timusk, T. and Statt, B. The pseudogap in high temperature superconductors: an experimental survey. *Reports of Progress in Physics*, to be published.
 - [21] Reznik, D. *et al.* Direct observation of optical magnons in $YBa_2Cu_3O_{6.2}$. *Phys. Rev. B*, **53** R14741-R14744 (1996)

- [22] Hayden, S. *et al.* High Frequency spin waves in $YBa_2Cu_3O_{6.15}$. *Phys. Rev. B*, **54** R6905-R6908 (1996)
- [23] Scalapino, D. The case for $d_{x^2-y^2}$ pairing in the cuprate superconductors. *Phys. Rep.* , **250** 329-365 (1995).
- [24] Pines, D. Understanding high-temperature superconductivity, a progress report. *Physica B*, **199-200** 300-309 (1994).
- [25] Shrieffer, J.R. Ward's identity and the suppression of spin fluctuation superconductivity. *J.Low Temp. Physics*, **99** 397-402 (1995).
- [26] Bulut, N. and Scalapino, D. Neutron scattering from a collective spin fluctuation mode in CuO_2 bilayer. *Phys. Rev. B*, **53** 5149-5152 (1996).
- [27] Morr, D.K. and Pines, D. The resonance peak in cuprate superconductors. *preprint*, cond-mat/9805107.
- [28] Hayden, S. *et al.* Absolute measurements of the high-frequency magnetic dynamics in high-Tc superconductors. *Physica B* **241-243** 765-772 (1998)
- [29] Fong, H.F. *et al.* Superconductivity-induced anomalies in the spin excitation spectra of underdoped $YBa_2Cu_3O_{6+x}$. *Phys. Rev. Lett.* **78** 713-716 (1997)
- [30] Loram, J.W. *et al.* Superconducting and normal state energy gaps in $Y_{0.8}Ca_{0.2}Ba_2Cu_3O_{7-\delta}$ from the electronic specific heat. *Physica C* **282-287** 1405-1406 (1997).
- [31] Dai, P. *et al.* Magnetic dynamics in undoped $YBa_2Cu_3O_{7-x}$: direct observation of a superconducting gap. *Phys. Rev. Lett.* **77** 5425-5428 (1996)